

## National Registry of Emergency Medical Technicians® Emergency Medical Responder Psychomotor Examination

## **CARDIAC ARREST MANAGEMENT / AED**

Candidate:	Examiner:		
Date:	Signature:		
Actual Time Started:		Possible Points	Points Awarded
Takes or verbalizes appropriate PPE precautions		1	
Determines the scene/situation is safe		1	
Attempts to question bystanders about arrest events		1	
Checks patient responsiveness		1	
Requests additional EMS assistance		1	
Checks breathing and pulse simultaneously		1	
NOTE: After checking responsiveness, then checking breathing examiner informs candidate, "The patient is unresponsive, apne		ds,	
Immediately begins chest compressions [adequate depth and rate; allo	ws the chest to recoil completely]	1	
Performs 2 minutes of high-quality, 1-rescuer adult CPR -Adequate depth and rate (1 point) -Correct compression-to-ventilation ratio (1 point) -Allows the chest to recoil completely (1 point) -Adequate volumes for each breath (1 point) -Minimal interruptions of no more than 10 seconds throughout (1 point)	int)	5	
NOTE: After 2 minutes (5 cycles), candidate assesses patient and second rescuer resumes compressions while candidate operates AED.			
Turns on power to AED		1	
Follows prompts and correctly attaches AED to patient		1	
Stops CPR and ensures all individuals are clear of the patient during rh	· · · · · · · · · · · · · · · · · · ·	1	
Ensures that all individuals are clear of the patient and delivers shock fi	rom AED	1	
Immediately directs rescuer to resume chest compressions		1	
Actual Time Ended:	TOTAL	17	
Critical Criteria  Failure to take or verbalize appropriate PPE precautions Failure to check responsiveness, then check breathing and pulse Failure to immediately begin chest compressions as soon as puls Failure to demonstrate acceptable high-quality, 1-rescuer adult Cl Interrupts CPR for more than 10 seconds at any point Failure to correctly attach the AED to the patient Failure to operate the AED properly Failure to deliver shock in a timely manner Failure to ensure that all individuals are clear of patient during rhy [verbalizes "All clear" and observes] Failure to immediately resume compressions after shock delivered Failure to manage the patient as a competent EMR Exhibits unacceptable affect with patient or other personnel Uses or orders a dangerous or inappropriate intervention	elessness is confirmed PR thm analysis and before delivering shock		

You must factually document your rationale for checking any of the above critical items on the reverse side of this form.